



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The status of the mixture is described in the table.

Professor Harry F. Reid informs me that a continental shelf of the Atlantic Ocean has a very definite slope which is very nearly that obtained for the extremely wet sand in these experiments.

I reserve the mathematical theory for a future paper.

SAND	WATER	ANGLE OF REPOSE	REMARKS
<i>pounds</i>	<i>pounds</i>		
10	0	33°	Dry
10	0.5	65°	Not hard
10	1.0	120°	Not accurate, but large obtuse angle, hard
10	1.5	120°-140°	Not accurate, but large obtuse angle, hard
10	2.0	120°-140°	Not accurate, but large obtuse angle, hard
10	2.5	120°	Not accurate, but large obtuse angle, hard
10	3.0	48°	Fairly hard
10	3.5	19°	All mixes
10	3.75	14.5°	Very slight excess of water
10	4.0	13°	Water not all absorbed
10	5.0	12°	Excess of water

The sand has been meshed by Professor Roys of the Worcester Polytechnic Institute with the following results:

SIZE OF SCREEN	DIAMETER OF OPENING	PER CENT OF SAND WHICH PASSED THROUGH
<i>meshes</i>	<i>inches</i>	
200	0.0029	7.3
100	0.0055	29.55
50	0.011	84.60
30	0.022	99.45
20	0.034	99.85
10	0.073	99.95
4	0.20	100.00

PALÆOMASTODON, THE ANCESTOR OF THE LONG-JAWED MASTODONS ONLY

BY HENRY FAIRFIELD OSBORN

AMERICAN MUSEUM OF NATURAL HISTORY, NEW YORK CITY

Read before the Academy, April 29, 1919

In 1900¹ the author predicted that the ancestors of the Proboscideans, as well as of the Hyracoidea and some other orders of mammals, would be discovered in Africa. Two years later the members of the British Geological Survey of Egypt discovered in the Oligocene of the Fayûm remains of *Palæo-*

mastodon and of *Mærittherium*, which were at once regarded as the solution of the ancestry of the Proboscideans. These animals took their place in all literature as two steps in the early evolution of this remarkable group.

In 1909² Osborn pointed out that *Mærittherium* is to be regarded as a terrestrial form of the Sirenians (manatees and dugongs) in no way directly related to the Proboscideans. It now appears that *Palæomastodon* must also be removed from its generalized position and be regarded as the ancestor of the long-jawed mastodons only; it is far too much specialized in the longirostral direction to be ancestral to the Proboscidea in general. These long-jawed mastodons are distinguished by the peculiar use of the front teeth of the lower jaw, which together made a broadly flattened, spoon-shaped tooth,




FIG. 1

Outline of the mounted skeleton of *Megabelodon* in The American Museum of Natural History, discovered by E. L. Troxell in the Middle Pliocene of Texas. Drawing one forty-fifth natural size.

almost entirely enamel covered. Phases of the evolution of this long-jawed phylum are seen in the classic *Trilophodon angustidens* of Cuvier, in the lower Miocene of France. A branch reached Texas in the Upper Miocene (*Trilophodon productus* of Cope), and Florida as well as Texas in the *Trilophodon serridens* of Cope. It attained gigantic proportions in the Middle Pliocene. The *Megabelodon* of Barbour, a superb skeleton of a long-jawed and extremely short-limbed Proboscidean, recently discovered in Texas by Mr. E. L. Troxell, has been mounted in The American Museum of Natural History. It represents one of the culminating stages in the evolution of the long-jawed mastodons. In these animals we find proof of nearly direct linear descent from the *Palæomastodon* of the Fayûm, e.g., the long enamel band on the upper tusks, the broadly spoon-shaped arrangement of the lower tusks with enamel covering. In massiveness these animals parallel and even surpass the true mastodons of the Pleistocene, to which they are only indirectly related.

¹Osborn, 1900, 182. ²Osborn, 1909, 332.